



Your comprehensive guide to

BREAST AUGMENTATION

Breast Augmentation

Breast augmentation, also known as augmentation mammoplasty, is a very popular and common procedure that can increase your breast size by one or more sizes. By surgically inserting breast implants beneath the breast tissue or the chest muscle during breast augmentation, women can achieve larger, shapelier breasts. This type of breast enhancement is sometimes performed in conjunction with a breast lift.

For some women, breast augmentation is a way to enhance self-image and self-confidence. For others, breast augmentation is part of breast reconstruction after surgery for breast cancer or other conditions affecting the breast.

Breast Augmentation Considerations:

Breast augmentation can dramatically improve the appearance of a woman's breasts. During your breast augmentation consultation, you'll learn more about your options for creating your ideal breast appearance, including:

- Implant size, style, and shape.
- Implant material (silicone gel, or saline).
- Implant position (above or below the pectoralis muscle).
- Implant incision site.

The goal is to create a natural and alluring look. Which implant size that will work best for you must be individually determined with your input. The size, shape, style, and type of implant will be decided after a comprehensive examination. Placing implants that are too large for the body size have greater chances of capsular contracture, discomfort and longer recovery times.

Breast Implant Incision Options

Inframammary incision:

This is the most commonly used incision at the present time and is made below the breast fold. This incision is generally well concealed and may cause fewer breastfeeding difficulties than the periareolar incision option.

Periareolar incision:

It is associated with a higher likelihood of breast feeding difficulties than the other incision options because periareolar incision involves cutting through the breast tissue. This may also increase the chance that there will be a change in nipple sensation.

Breast Implant Placement Options

Sub-muscular implant placement:

The breast implants are placed partially or wholly under the pectoralis major chest muscle. Sub-muscular implants may result in a slightly longer, more painful recovery than sub-glandular implants. Sub-muscular placement may result in less palpable implants, as well as offer easier mammographic imaging of the breast. One potential disadvantage is implant movement with muscle action and so can be an issue with the gym.

Sub-glandular implant placement:

The breast implant location is between your chest muscle and your breast tissue (under the glands). This placement may help reduce your surgery and recovery time. It may be less painful than sub-muscular placement and may make your breast implants easier to access if re-operation is necessary. Implants that are sub-glandular may be easier to see and feel through your skin.

Each placement type has benefits that may be right for you. During your consultation you will discuss your desired outcome with Dr John Flynn to select the best option for your figure.

Procedure – What to Expect

Breast augmentation can be performed in a surgical centre or hospital outpatient facility. Surgery usually takes one to two hours, and is performed under general anaesthesia. You will expect to go home the same day of your surgery (rarely does the procedure require a hospital stay).

Dr John Flynn will give you specific guidelines to follow during your recovery based on your personal needs. In general, you can expect to return to your usual routine after a few days, but you should avoid excessive lifting to minimise tension on incision lines, thereby ensuring minimal scar lines.

Recovery – What to Expect

Feeling tired and mild soreness following breast augmentation surgery are common. Pain medication and antibiotics may be prescribed to help alleviate discomfort and reduce the risk of infection.

After breast augmentation surgery you may feel a burning sensation in your nipples that will fade after about two weeks. It is advised that you remain in a support bra with no underwires for approximately four to six weeks. Breast swelling may take up to five weeks to subside. We recommend no gym or jogging for 4-6 weeks.

Scars usually stay firm and pink for at least six weeks and will begin to fade after several months, taking on a flesh tone by 6 to 9 months post-operatively.

Breast Augmentation Surgery Risks

Each year thousands of women undergo successful breast augmentation surgery, experience no major problems and are happy

with the results. Significant complications from a breast augmentation are infrequent. However, make sure you understand what surgery involves, including possible risks, complications and follow-up care.

Breast augmentation poses various risks, including:

- Capsular contracture (a tightening of the scar tissue which can make the breast feel hard and can distort the shape of the breast implant).
- Changes in nipple and breast sensation (usually temporary).
- Breast pain.
- Rare implant leakage or rupture. If the implant is saline-filled, it deflates and the body harmlessly absorbs the saline solution. If the implant is filled with silicone gel, the gel may alter the shape of the breast, requiring implant replacement.

Correcting any of these complications might require additional surgery, either to remove or replace the implants.

Other Considerations for breast augmentation surgery include:

- Breast implants won't prevent your breasts from sagging. A breast lift may be needed in addition to a breast augmentation to correct sagging breasts.
- Breast implants aren't guaranteed to last a lifetime. In addition to the possibility of implant rupture or leakage, factors such as aging, weight gain and/or weight loss may further change the way your breasts look. Any of these issues might lead to additional surgery.
- Mammograms might be more complicated. If you have breast implants, routine mammograms might require additional, specialised views or ultrasound.
- Breast implants might hamper breast-feeding. Most women are able to successfully breastfeed after breast augmentation. For others, however, breastfeeding is a challenge with or without breast implants.
- Insurance might not cover breast implants. Unless it's medically necessary, such as the need for implant reconstruction after a mastectomy, breast augmentation is not covered by insurance. Be prepared to handle any expenses that accompany breast augmentation, including related surgeries or future imaging tests.
- Additional surgery may be needed after breast implant removal. If you decide to have your implants removed at a later time in your life, you might need a breast lift or other corrective surgery to help your breasts maintain an aesthetically pleasing appearance.

Like any major surgery, breast augmentation surgery poses a risk of bleeding, infection and an adverse reaction to anaesthesia. It's also possible to have an allergic reaction to the surgical tape or other materials used during or after the procedure.

You can help minimise certain risks by following the advice and instructions of Dr Flynn, both before and after your breast augmentation surgery. If you have any concerns about the risks involved with breast augmentation surgery, please consult Dr Flynn or his team at Cosmedic & Skin.

Breast Augmentation Cost

Breast augmentation by Dr John Flynn on the Gold Coast starts from \$11,990. The cost for breast surgery may vary based on your individual requirements and surgical desires. The exact cost for surgery will be provided during your initial consultation.



Contact Us

(07) 5588 4777

cosmedic.com.au

Pivotal Point, Suite 203, Level 2

50 Marine Parade, QLD 4215